

Usability

- ONC's Unified Understanding
 - and the
 - Value of SHARP C

Joe Bormel, MD, MPH Tuesday, March 25, 2014





ADVERSITY

THAT WHICH DOES NOT KILL ME POSTPONES THE INEVITABLE.

Abdominal Pain



Share this:









Just about everybody at one point or another will experience abdominal pain. Most of the causes are not serious and can be readily diagnosed and treated. However, pain can also be a sign of a serious illness. It's important to be able to recognize symptoms that are severe and know when to call a doctor.

What Are the Most Common Causes of Abdominal Pain?

Whether it's a mild stomach ache, sharp pain, or stomach cramps, abdominal pain can have numerous causes. Some of the more common causes include:

- Indigestion
- Constipation
- Stomach virus
- Menstrual cramps
- Food poisoning
- Food allergies
- Gas
- Lactose intolerance
- Ulcers
- Pelvic inflammatory disease
- Hernia
- Gallstones
- Kidney stones
- Endometriosis









@Mayo Foundation for Medical Education and Research. All rights reserved.

Usability travels with Safety



Information Access Division (IAD)



What is Usability?

Usability is "the extent to which a product can be used by specified users to achieve specified goals with effectiveness, efficiency and satisfaction in a specified context of use"

- ISO 9241

Visualization and Usability Group: Ilowry@nist.gov

The Two Bins of Usability

User Experience (UX Design)

User Interface Design

Displays and Controls
Screen Design
Clicks & Drags


Cognitive Task Support


"Workflow Design"
Data Visualization
Functionality


Usability and User Centered Design

 Usability: Extent to which a product can be used by <u>specified</u> users to achieve <u>specified goals</u> with <u>effectiveness</u>, <u>efficiency</u> and <u>satisfaction</u> in a <u>specified context</u> of use

• UCD









A MORE REALISTIC CHESS

Because the world is not just black and white

Rather than being the main instigators of an accident, operators tend to be the inheritors of system defects..... Their part is that of adding the final garnish to a lethal brew that has been long in the cooking."

James Reason, Human Error, 1990

Standard PCA Parameters for Opioid Naïve Adult Patients					
	morphine	HYDROmorphone	fentaNYL		
1x (single strength)	1mg/ml	0.2 mg/ml (200 mcg/ml)	10 mcg/ml		
Loading Bolus	2 mg	0.4 mg (400 mcg)	20 mcg		
Clinician Bolus	2 mg	0.4 mg (400 mcg)	20 mcg		
Number of Clinician Boluses Per Hour	1	1	1		
PCA Dose	1 mg	0.2 mg (200 mcg)	10 mcg		
Lockout	10 minutes	10 minutes	10 minutes		
Total Drug Over Time	Optional	Optional	Optional		
Max Number of Patient Demand Doses Per Hour	Optional	Optional	Optional		
Basal	Not recommended for starting PCA				


What creates or destroys Usability?











Workflow

- Workflow modeling Methods
- http://nvlpubs.nist.gov/n istpubs/ir/2014/NIST.IR. 7988.pdf


Integrating Electronic Health Records into Clinical Workflow: An Application of Human Factors Modeling Methods to Ambulatory Care

Sventlana Z. Lowry
Mala Ramaiah
Emily S. Patterson
David Brick
Ayse P. Gurses
Ant Ozok
Debora Simmons
Michael C. Gibbons

http://dx.doi.org/10.6028/NIST.IR.7988







Know Your Brain, Transform Your Performance


STRATEGIES FOR OVERCOMING DISTRACTION, REGAINING FOCUS, & WORKING SMARTER ALL DAY LONG

David Rock

Foreword by Daniel J. Siegel, M.D.

• Think of conscious thinking as a precious resource Some Things to Try

- Prioritize prioritizing, as it's an energy-intensive Save mental energy for prioritizing by avoiding
 - other high-energy-consuming conscious activities • Schedule the most attention-rich tasks when you
 - Use the brain to interact with information rather
 - than trying to store information, by creating visuals • Schedule blocks of time for different modes of
 - thinking.







THINKING,
FAST AND SLOW



DANIEL KAHNEMAN

WINNER OF THE NOBEL PRIZE IN ECONOMICS





PERSPECTIVE

Just remember, whatever side you're on the other person is right too.

"A landrark contribution in humanity's understanding of Burd."

— The New York Down Book Senters

THE RIGHTEOUS MIND

WHY GOOD
PEOPLE ARE DIVIDED
BY POLITICS AND
RELIGION

JONATHAN HAIDT



"Your mother and I are separating because I want what's best for the country and your mother doesn't."

FIGURE 13.1. Why Manichaeans think they are divided by politics. (photo credit 13.1)

People don't adopt their ideologies at random, or by soaking up whatever ideas are around them. People whose genes gave them brains that get a special pleasure from novelty, variety, and diversity, while simultaneously being less sensitive to signs of threat, are predisposed





FIGURE 6.1. Two dimensions of cognitive style. People with autism are very high on systemizing and very low on empathizing. So were some important moral philosophers. (Adapted from Baron-Cohen 2009.)

Flavor of the Frustration

- "Every Screen's got 50 different things, you know, that are changing. It slows me down. So, I do a lot of my charting at night...But, the problem is I'm spending more hours doing it than I would have before. We have XXX, which you have to be careful of, because I just [dictated] a 'Patient's prostate is bothering him' and it turned out 'Patient's prostitute is bothering him.' You really have to read careful, because I can end up going to court with that stuff."
 - Primary care physician








MAGIC

Usability Is ... Personal



Usability Requires Experimentation


Lesson from Twinlist: a little bit of visualization can go a long way!

From: Krisanne Graves [mailto:Krisanne.Graves@uth.tmc.edu] Sent: Friday, November 22, 2013 11:23 AM To: 'Krisanne Graves'; Amy Franklin; Dean F. Sittig; Jiajie Zhang; Trevor Cohen; Jorge R. Herskovic; Carmen Elliott; Hua Xu; Ada (awright5@partners.org); Ben Shneiderman (ben@cs.umd.edu); Catherine Plaisant (plaisant@cs.umd.edu); Keith Butler (keith.a.butler@gmail.com); Robert A. Greenes (greenes@asu.edu); Reider, Jacob (HHS/ONCIT); Morton, Alicia (HHS/ONC); Di Sottara; Charles, Dustin (HHS/ONC); Belden, Jeffery L.; Muhammad F. Walji - UT Houston; Helwig, Amy (HHS/ONCIT); Lana Lo (svetlana.lowry@nist.gov); Bormel, Joseph (HHS/ONCIT); AHRQ Johnson, Todd Cc: Pettis, Luciana Subject: SHARPC Project Leaders Meeting When: Occurs the first Monday of every 1 month effective 1/6/2014 until 3/31/2014 from 2:00 PM to 4:00 PM Central Standar Where: Webex and Teleconference TBA

Location: Webex and Teleconference TBA Account ' When: Occurs the first Monday of every Bormel, Joseph (HHS/ONCIT) on be From: Adam Wright (awright5 @partners.org) 1 2 AHRQ Todd Johnson TAB Alicia Morton (HHS/ONC) Amy Franklin Amy Helwig (HHS/ONCIT) 6 Ben Shneiderman (ben @cs.umd.edu) 7 Carmen Elliott Catherine Plaisant (plaisant @cs.umd.edu) 9 Davide Sottara 10 Dean F. Sittig 11 **Dustin Charles** (HHS/ONC) 12 Hua Xu 13 Jacob Reider-----(HHS/ONCIT) 14 Jeffery L Belden TAB 15 Jiaije Zhang 16 Jorge, R Herskovic 17 Joseph Bormel (HHS/ONCIT) 18 Keith Butler (keith a butler @gmail.com) 19 Krisanne Graves 20 Lana Lowry (svetlana lowry @nist.gov) 21 Luciana Pettis TAB 22 Muhammad F Walii _ UT Houston 23 Robert A Greenes (greenes @asu.edu) 24 Trevor Cohen

Usability Requires Learning

Research/Experimentation, Prototyping, Testbeds, Scenarios



Article Talk Read Edit View his

Self-experimentation


From Wikipedia, the free encyclopedia

Self-experimentation refers to the very special case of single-subject *scientific* experimentation experiment on her- or himself. Usually this means that the designer, operator, subject, analyst, an the same. It is a special case of single-subject research.



	PROTOTYPING	LIVE PROTOTYPING	TECHNICAL PROTOTYPING	PILOT
Question answered	What are the different ways we could solve this problem?	Does our message and solution resonate in the market?	Can the problem be solved in this specific way?	Are the economic attractive enough to justify scaling?
When to employ it in the product- development process	Early, to explore many ways to achieve a goal and periodically during the midphase to explore ways to deliver a specific feature or value driver.	Midway, to rigorously explore market appeal once a specific opportunity area has been identified.	When technical feasibility is in question or when the function has been identified but the specific means is yet to be determined.	Late, as a final check to tweak details before launching at scale
Fidelity (i.e. the degree to which a prototype reflects a polished and finished product)	Low	Just enough for the product to appear real in the marketplace.	High enough to prove feasibility.	Very high. Pilots require a working solution and are often more polished than the scalable versions to overcome initial market awarenes challenges.


Learning Requires Liberty



Adaptive Checklists









Optimum Strategic Position for ONC to Leverage its Regulatory Authority for Change Health IT Po




the National Coordinator for Health IT



The Learning Healthcare System



Usability in Healthcare



Health eDecisions (HeD)

www.healthedecisions.org

- Goal:
- To define and validate standards
- that facilitate the emergence of systems and services whereby CDS interventions
- can be shared or accessed
- by any healthcare stakeholder
- via an importable format or via a CDS Web service.
 - Outcome: → enable CDS sharing at scale

EHR	Pilot	Content Supplier
Design Clinicals	Order Set – Heart Failure	Zynx Health
Allscripts	Rule – NQF 068 (Million Hearts)	newMentor
Allscripts	Rule – San Diego Pertussis	CDC
VA	Documentation Template – UTI	Wolters Kluwer Health

Knowicage and the


Learning Health System



Standardized Knowledge



Table 1: Standard PCA Parameters for Opioid <u>Naïve</u> Adult Patients				
Standard PCA Parameters for Opioid <u>Naïve</u> Adult Patients				
	morphine	HYDROm orphone	fenta NY L	
1x (single strength)	1mg/ml	0.2 mg/ml (200 mcg/ml)	10 mcg/ml	
Loading Bolus	2 mg	0.4 mg (400 mcg)	20 mcg	
Clinician Bolus	2 mg	0.4 mg (400 mcg)	20 mcg	
Number of Clinician Boluses Per Hour	1	1	1	
P CA Dose	1 mg	0.2 mg (200 mcg)	10 mcg	
Lockout	10 minutes	10 minutes	10 minutes	
Total Drug Over Time	Optional	Optional	Optional	
Max Number of Patient Demand Doses Per Hour	Optional	Optional	Optional	
Basal	Not recommended for starting P CA			




Improved & Validated Knowledge



People

Process





Skills

Improved & Validated
Knowledge

Tools

Workplace

Thank You

- Joe Bormel, MD, MPH
- ONC Office of the Chief Medical Officer
 - Joseph.Bormel@hhs.gov